arXiv:1401.3708v1 [math.DS] 15 Jan 2014

CLASSIFYING $GL(2,\mathbb{Z}) \ltimes \mathbb{Z}^2$-ORBITS BY SUBGROUPS OF $\mathbb{R}$

DANIELE MUNDICI

Abstract. Let $\mathcal{G}_2$ denote the affine group $GL(2,\mathbb{Z}) \ltimes \mathbb{Z}^2$. For every point $x = (x_1, x_2) \in \mathbb{R}^2$ let $\mathrm{orb}(x) = \{y \in \mathbb{R}^2 \mid y = \gamma(x) \text{ for some } \gamma \in \mathcal{G}_2\}$. Let G_x be the subgroup of the additive group $\mathbb{R}$ generated by $x_1, x_2, 1$. If $\mathrm{rank}(G_x) \in \{1, 3\}$ then $\mathrm{orb}(x) = \{y \in \mathbb{R}^2 \mid G_y = G_x\}$. If $\mathrm{rank}(G_x) = 2$, knowledge of G_x is not sufficient in general to uniquely recover $\mathrm{orb}(x)$: rather, G_x classifies precisely $\max(1, \phi(d)/2)$ different orbits, where d is the denominator of the smallest positive nonzero rational in G_x and ϕ is Euler function. To get a complete classification, polyhedral geometry provides an integer $c_x \geq 1$ such that $\mathrm{orb}(y) = \mathrm{orb}(x)$ iff $(G_x, c_x) = (G_y, c_y)$.

1. Introduction

Throughout we let $\mathcal{G}_n = GL(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$ denote the group of affine transformations of the form $x \mapsto Ux + t$ for $x \in \mathbb{R}^n$, where $t \in \mathbb{Z}^n$ and U is an integer $n \times n$ matrix with $\det(U) = \pm 1$. By an "orbit" we will mean a $\mathcal{G}_n$-orbit. The dimension of the ambient space $\mathbb{R}^n$ will always be clear from the context. We let $\mathrm{orb}(x)$ denote the orbit of $x \in \mathbb{R}^n$. Thus

$$\mathrm{orb}(x) = \{y \in \mathbb{R}^n \mid y = \gamma(x) \text{ for some } \gamma \in \mathcal{G}_n\}.$$

This paper is devoted to the classification of $\mathcal{G}_2$-orbits: to every point in $\mathbb{R}^2$ we will assign an invariant, in such a way that two points have the same orbit iff they have the same invariant. For each $x = (x_1, \ldots, x_n) \in \mathbb{R}^n$, the main invariant considered in this paper is the group

$$G_x = \mathbb{Z} + \mathbb{Z}x_1 + \mathbb{Z}x_2 + \cdots + \mathbb{Z}x_n$$

generated by $1, x_1, x_2, \ldots, x_n$ in the additive group $\mathbb{R}$. In Propositions 2.1 and 2.3 (also see Remark 2.4) it is shown that G_x completely classifies the orbit of any $x \in \mathbb{R}^n$, $n \geq 2$, whenever $\mathrm{rank}(G_x)$ equals 1 or $n + 1$.

If $x \in \mathbb{R}^2$ and $\mathrm{rank}(G_x) = 2$, the invariant G_x is not complete for $\mathrm{orb}(x)$. Rather, G_x classifies precisely $\max(1, \phi(d)/2)$ different orbits, where d is the denominator of the smallest positive nonzero rational in G_x and ϕ is Euler function. To get a complete invariant we assign to x an integer $c_x \geq 1$ and prove in Theorem 6.1 that $\mathrm{orb}(x) = \mathrm{orb}(y)$ iff $(G_x, c_x) = (G_y, c_y)$. In Corollary 6.3 we describe the totality of pairs (G_x, c_x) arising as complete classifiers of the totality of orbits of points of $\mathbb{R}^2$. The geometric significance of c_x is discussed in Section 5.

Date: January 16, 2014.

Key words and phrases. Affine group over the integers, orbit, $GL(2,\mathbb{Z}) \ltimes \mathbb{Z}^2$-orbit, complete invariant, rational simplex, regular cone, Farey sequence, totally ordered abelian group, AF C*-algebra.

2000 *Mathematics Subject Classification.* Primary: 37C85. Secondary: 11B57, 22F05, 37A45, 06F20, 46L80.

While the study of the orbits of $\mathcal{G}_2$ and of its subgroups $GL(2,\mathbb{Z})$, $SL(2,\mathbb{Z})$, $SL(2,\mathbb{Z})_+$ requires techniques from various mathematical areas [3, 8, 10, 15, 16, 17], a first possible reason of interest in our classification stems from the novel role played by polyhedral geometry [6], through the fundamental notion of regular cone and regular simplex. (As is well known, a regular fan is a complex of regular cones.) Secondly and thirdly, in a final section we will apply our classification to the analysis of germinal ideals of lattice-ordered abelian groups with strong order unit [1],[13] and of the Farey-Stern-Brocot AF C*-algebra, [12, 2, 4, 14].

2. Rank 3 and Rank 1

Fix $n = 1, 2, \ldots$. In case $x \in \mathbb{R}^n$ and $\mathrm{rank}(G_x) = n+1$, the following elementary result shows that the group G_x is a complete classifier of the orbit of x. Its proof is left to the reader.

Proposition 2.1. *Let us say that a point $x = (x_1, \ldots, x_n) \in \mathbb{R}^n$ is* totally irrational *if the $(n+1)$-tuple $(x_1, \ldots, x_n, 1)$ is rationally independent. In other words,* $\mathrm{rank}(G_x) = n+1$. *Whenever x is totally irrational,* $\mathrm{orb}(x)$ *is completely classified by G_x. Thus in particular, if $x \in \mathbb{R}^2$ and* $\mathrm{rank}(G_x) = 3$ *then* $\mathrm{orb}(x) = \{y \in \mathbb{R}^2 \mid G_y = G_x\}$.

To deal with the remaining cases we introduce some basic tools from polyhedral geometry [6], [13], featuring the notion of regular cone and simplex.

Rational points, simplexes and cones. A point $y = (y_1, \ldots, y_n) \in \mathbb{R}^n$ is said to be *rational* if $y_i \in \mathbb{Q}$ for each $i = 1, \ldots, n$. Equivalently, $\mathrm{rank}(G_x) = 1$. By the *denominator* $\mathrm{den}(y)$ of the rational point $y = (y_1, \ldots, y_n)$ we understand the least common denominator of its coordinates. Observe that, for each rational point $y \in \mathbb{R}^n$, G_y is the group of all integer multiples of $1/\mathrm{den}(y)$. A simplex $T \subseteq \mathbb{R}^n$ is said to be *rational* if each vertex of T is a rational point. A *rational hyperplane* $H \subseteq \mathbb{R}^n$ is a set of the form $H = \{z \in \mathbb{R}^n \mid h \circ z = k\}$, for some nonzero vector $h \in \mathbb{Q}^n$ and $k \in \mathbb{Q}$. A *rational affine space* A in $\mathbb{R}^n$ is the intersection of rational hyperplanes in $\mathbb{R}^n$. If the origin belongs to A we say that A is *homogeneous*.

Following [6, p.6], given vectors $v_1, \ldots, v_s \in \mathbb{R}^n$ we write

$$\mathrm{pos}[v_1, \ldots, v_s] = \mathbb{R}_{\geq 0}\, v_1 + \cdots + \mathbb{R}_{\geq 0}\, v_s$$

for their *positive hull* in $\mathbb{R}^n$. A nonzero integer vector $v \in \mathbb{Z}^n$ is said to be *primitive* ("visible" in [9]) if it is minimal, as a nonzero integer vector, along its *ray* $\mathrm{pos}[v]$. Further, [6, p.146], for $t = 1, 2, \ldots, n$, a *t-dimensional rational simplicial cone* in $\mathbb{R}^n$ is a set $C \subseteq \mathbb{R}^n$ of the form $C = \mathrm{pos}[w_1, \ldots, w_t]$, for linearly independent primitive integer vectors $w_1, \ldots, w_t \in \mathbb{Z}^n$. The vectors $w_1, \ldots, w_t$ are called the *primitive generating vectors* of C. They are uniquely determined by C. By a *face* of C we mean the positive hull of a subset of $\{w_1, \ldots, w_t\}$. For the sake of completeness we stipulate that the face of C determined by the empty set is the singleton $\{0\}$. This is the only zerodimensional cone in $\mathbb{R}^n$. By definition, the *homogeneous correspondent* of y is the primitive integer vector

$$\tilde{y} = (\mathrm{den}(y) \cdot y_1, \ldots, \mathrm{den}(y) \cdot y_n,\ \mathrm{den}(y)) = \mathrm{den}(y)(y, 1) \in \mathbb{Z}^{n+1}.$$

Given $k = 0, 1, \ldots$ and a rational k-simplex $T = \mathrm{conv}(v_0, \ldots, v_k) \subseteq \mathbb{R}^n$, we denote by $T^{\uparrow}$ the positive hull in $\mathbb{R}^{n+1}$ of the homogeneous correspondents of the vertices

of T. In symbols,

$$T^{\uparrow} = \mathrm{pos}[\tilde{v}_0, \ldots, \tilde{v}_k] = \mathbb{R}_{\geq 0}\tilde{v}_0 + \cdots + \mathbb{R}_{\geq 0}\tilde{v}_k \ \subseteq \mathbb{R}^{n+1}.$$

We say that $T^{\uparrow}$ is the *(rational simplicial) cone of* T. Note that $\dim(T^{\uparrow}) = k+1$.

Regularity. In rational polyhedral topology the following notion is fundamental, [6, p.146-147]:

Let $m = 1, 2, \ldots$ and $t = 1, \ldots, m$. Then a t-dimensional rational simplicial cone $C = \mathrm{pos}[d_1, \ldots, d_t] \subseteq \mathbb{R}^m$ is said to be *regular* if the set $\{d_1, \ldots, d_t\}$ of its primitive generating vectors can be extended to a basis of the free abelian group $\mathbb{Z}^m$ of integer points in $\mathbb{R}^m$. A rational simplex $T \subseteq \mathbb{R}^n$ is said to be *regular* (*unimodular* in [13]) if so is its associated cone $T^{\uparrow} \subseteq \mathbb{R}^{n+1}$.

The following elementary lemma will be our key tool to construct maps $\gamma \in \mathcal{G}_n$:

Lemma 2.2. *Let* $T = \mathrm{conv}(v_1, \ldots, v_{n+1})$ *and* $T' = \mathrm{conv}(v'_1, \ldots, v'_{n+1})$ *be regular* n*-simplexes in* $\mathbb{R}^n$*, with* $\mathrm{den}(v_i) = \mathrm{den}(v'_i)$ *for all* $i = 1, \ldots, n+1$*. Then for a unique* $\gamma \in \mathcal{G}_n$ *we have* $\gamma(v_1) = v'_1, \ldots, \gamma(v_{n+1}) = v'_{n+1}$.

Proof. The vectors $\tilde{v}_1, \ldots, \tilde{v}_{n+1}$ constitute a basis of the free abelian group $\mathbb{Z}^{n+1}$, and so do the vectors $\tilde{v}'_1, \ldots, \tilde{v}'_{n+1}$. Let V be the $(n+1) \times (n+1)$ matrix whose columns are $\tilde{v}_1, \ldots, \tilde{v}_{n+1}$. Let similarly V' be the matrix with columns $\tilde{v}'_1, \ldots, \tilde{v}'_{n+1}$. There is a unique (automatically integer and unimodular) $(n+1) \times (n+1)$ matrix M such that $MV = V'$. The assumption $\mathrm{den}(v_i) = \mathrm{den}(v'_i)$ for all i, is to the effect that the bottom row of M has the form $0, \ldots, 0, 1$ with n zeros. From M one immediately obtains the desired $\gamma \in \mathcal{G}_n$. □

For each $d = 1, 2, \ldots$, in the notation of [9, 3.1], we let $\mathfrak{F}_d$ be the dth *Farey sequence*, the ordered list of all rational numbers in $[0, 1]$ of denominator $\leq d$.

We are now in a position to show that G_x is a complete invariant for orbits of points $x \in \mathbb{R}^2$ with $\mathrm{rank}(G_x) = 1$:

Proposition 2.3. *If* $x \in \mathbb{R}^2$ *and* $\mathrm{rank}(G_x) = 1$ *then* $\mathrm{orb}(x) = \{x' \in \mathbb{R}^2 \mid G_{x'} = G_x\}$.

Proof. Trivially, $\mathrm{orb}(x) \subseteq \{x' \in \mathbb{R}^2 \mid G_{x'} = G_x\}$. For the converse inclusion, let $d = \mathrm{den}(x)$. Then $G_x = \mathbb{Z}\frac{1}{d}$. If $x' \in \mathbb{R}^2$ satisfies $G_{x'} = G_x$ then x' is a rational point and $\mathrm{den}(x') = d$. If $d = 1$ a translation by an integer vector will do. If $d > 1$ it is no loss of generality to assume that x and x' both lie in the unit square $[0, 1]^2$. On the ray $\mathbb{R}_{\geq 0}x$, let v be the nonzero integer vector nearest to the origin. Let O, e_1, e_2 respectively denote the origin in $\mathbb{R}^2$, the point $(1, 0)$ and the point $(0, 1)$. Then, as is well known, some affine transformation $\alpha \in GL(2, \mathbb{Z})$ satisfies $\alpha(v) = e_1$. As a consequence, α sends x to some rational point $r = (p/d, 0)$ lying on the unit interval $\mathrm{conv}(O, e_1)$. Similarly, some affine transformation $\beta \in GL(2, \mathbb{Z})$ sends x' to some point $r' = (p'/d, 0) \in \mathrm{conv}(O, e_1)$. Let q/c and s/t be the two neighbors (=contiguous elements) of r in the Farey sequence $\mathfrak{F}_d$, recalling that $d > 1$. Both denominators c and t are strictly smaller than d. If $d \geq 3$, then $c \neq t$ and we may assume $c < t$. If $d = 2$ we have $c = t = 1$. Let the integer vector $w \in \mathbb{R}^2$ be defined by $w = (q, 1 - c)$. Then the triangle $T = \mathrm{conv}(r, w, e_2)$ is regular: this is a consequence of the unimodularity property of any two neighbors in a Farey sequence. The denominators of r, w, e_2 are $d, 1, 1$, respectively. Letting similarly the triangle $T' = \mathrm{conv}(r', w', e_2)$ be defined with reference to r', it follows that the

denominators of the vertices r', w', e_2 of T' are again $d, 1, 1$. By Lemma 2.2 there is a map $\gamma \in \mathcal{G}_2$ satisfying $\gamma(r) = r'$, $\gamma(w) = w'$, $\gamma(e_2) = e_2$. Letting $\delta \in \mathcal{G}_2$ be defined by composition as $\delta = \beta^{-1}\gamma\alpha$, we conclude that $\delta(x) = x'$, as desired. □

Remark 2.4. The generalization of Proposition 2.3 to any dimension $n \geq 2$ is straightforward: for a proof of the nontrivial direction, for every $x \in \mathbb{Q}^n$ we can assume $\mathrm{den}(x) = d > 1$ (otherwise the result is trivial), and write $x = (p/d, 0, 0, \ldots, 0)$ for a suitable integer $0 < p < d$ with $\gcd(p, d) = 1$. Let F be the subspace of $\mathbb{R}^n$ spanned by x and e_2, where $e_2 = (0, 1, 0, \ldots, 0)$. As in the proof of Proposition 2.3, for some integer vector $w \in F$ there is a regular 2-simplex $T = \mathrm{conv}(x, e_2, w)$ with $\mathrm{den}(w) = 1$. A routine construction now represents T as a face of some regular n-simplex, $U = \mathrm{conv}(x, e_2, w, v_3, \ldots, v_n)$ in $\mathbb{R}^n$, for suitable integer vectors $v_3, \ldots, v_n$. Similarly, given $x' \in \mathbb{Q}^n$, the hypothesis $G_{x'} = G_x$ means that $\mathrm{den}(x') = \mathrm{den}(x) = d$. Without loss of generality we can write $x' = (p'/d, 0, 0, \ldots, 0)$, and show that also x' is the vertex of some regular n-simplex U' in $\mathbb{R}^n$ whose remaining vertices are integer vectors. By Lemma 2.2 some $\gamma \in \mathcal{G}_n$ satisfies $\gamma(x) = x'$.

The case $n = 1$ will be dealt with in Corollary 6.4.

3. Rank 2: More on regularity

The classification of orbits of points $x \in \mathbb{R}^2$ with $\mathrm{rank}(G_x) = 2$ is more laborious: the following example shows that in general, G_x alone is not a complete classifier of $\mathrm{orb}(x)$:

Example 3.1. *Let $a = (1/5, \xi)$ and $b = (2/5, \xi)$, for some irrational $\xi \in [0, 1]$. Then $G_a = G_b$ but for no $\gamma \in \mathcal{G}_2$ we can have $\gamma(a) = b$.*

Proof. For otherwise (absurdum hypothesis), we can write

$$\begin{pmatrix} x & y & z \\ u & v & w \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1/5 \\ \xi \\ 1 \end{pmatrix} = \begin{pmatrix} 2/5 \\ \xi \\ 1 \end{pmatrix}$$

for some unimodular integer 3×3 matrix U. Since ξ is irrational, from $x/5 + y\xi + z = 2/5$ we get $y = 0$ and $x = 2 - 5z$. Further, from $u/5 + v\xi + w = \xi$ we get $v = 1$ and $w = -u/5$. In conclusion, the unimodularity of U yields

$$\det \begin{pmatrix} 2 - 5z & 0 \\ u & 1 \end{pmatrix} = \pm 1,$$

whence $2 - 5z = \pm 1$, which is impossible. □

In Definition 5.4 we will introduce a second invariant $c_x \in \mathbb{Z}$ and show that the pair (G_x, c_x) provides a complete classification of $\mathcal{G}_2$-orbits.

For later purposes we record here the following easy lemma:

Lemma 3.2. *For $n = 1, 2, \ldots$, let F be an e-dimensional rational affine space in $\mathbb{R}^n$ with $0 \leq e < n$. Let $\mathrm{conv}(v_0, \ldots, v_e)$ and $\mathrm{conv}(w_0, \ldots, w_e)$ be two regular e-simplexes lying on F. Suppose $\mathrm{conv}(v_0, \ldots, v_e, v_{e+1}, \ldots, v_n)$ is a regular n-simplex, for suitable points $v_{e+1}, \ldots, v_n \in \mathbb{R}^n$. Then also $\mathrm{conv}(w_0, \ldots, w_e, v_{e+1}, \ldots, v_n)$ is a regular n-simplex.*

The following characterization of regularity will find repeated use throughout the rest of this paper:

Lemma 3.3. *Given integers $n \geq 1$ and and $0 \leq m \leq n$, let $T = \mathrm{conv}(v_0, \ldots, v_m)$ be an m-simplex in $\mathbb{R}^n$. Then the following conditions are equivalent:*

(i) *T is regular.*

(ii) *The half-open parallelepiped $P_T = \{x \in \mathbb{R}^{n+1} \mid x = \sum_{i=0}^{m} \lambda_i \tilde{v}_i,\ 0 \leq \lambda_i < 1\}$ contains no nonzero integer point.*

(iii) *For every nonempty face F of T and every rational point r lying in the relative interior of F, the denominator of r is $\geq$ the sum of the denominators of the vertices of F.*

Proof. (i)⇒(ii) Trivial. (ii)⇒(i) If T is not regular some point q of the abelian group $\mathbb{Z}^{n+1} \cap (\mathbb{R}\,\tilde{v}_0 + \cdots + \mathbb{R}\,\tilde{v}_m)$ is not expressible as a linear combination of $\tilde{v}_0, \ldots, \tilde{v}_m$ with integer coefficients. From q one easily obtains by translation a nonzero integer point q' lying in P_T.

For the proof of (iii)⇔(i), supposing $W = \mathrm{conv}(z_1, \ldots, z_k)$ is a nonempty face of T we will write $g_i = \tilde{z}_i$ $(i = 1, \ldots, k)$ for the primitive generating vectors of the cone $W^{\uparrow} = \mathrm{pos}[g_1, \ldots, g_k]$, and P_W for the half-open parallelepiped associated to W. We further let Σ_W denote the sum of the denominators of the vertices of W. In more detail, letting $g_{i,n+1}$ denote the $(n+1)$th coordinate of g_i of $W^{\uparrow}$, $\Sigma_W = g_{1,n+1} + g_{2,n+1} + \cdots + g_{k,n+1}$.

(iii)⇒(i) If T is not regular, by (ii)⇒(i) the half-open parallelepiped P_T contains a nonzero integer point $u = (u_1, \ldots, u_{n+1}) \in \mathbb{Z}^{n+1}$. Without loss of generality, u is primitive. Let a be the uniquely determined rational point such that $u = \tilde{a}$. It follows that $a \in T$ and $\mathrm{den}(a) = u_{n+1} = (n+1)$th coordinate of u. Let S be the smallest face of T such that a belongs to S. Then a lies in the relative interior of S, and u belongs to the half-open parallelepiped P_S. Therefore, $\mathrm{den}(a) < \Sigma_S$ and condition (iii) fails.

(i)⇒(iii) Suppose T is regular, $F = \mathrm{conv}(w_0, \ldots, w_l)$ is a nonempty face of T, and r is a rational point in the relative interior of F. Then F is regular, and the homogeneous correspondent $\tilde{r}$ of r is a primitive integer vector lying in the relative interior of the cone $F^{\uparrow}$. For suitable integers $n_i \geq 1$ we can write $\tilde{r} = n_0\tilde{w}_0 + \cdots + n_l\tilde{w}_l$, whence the $(n+1)$th coordinate $\tilde{r}_{n+1} = \mathrm{den}(r)$ of $\tilde{r}$ satisfies the inequality $\tilde{r}_{n+1} \geq \Sigma_F$. □

4. Rank 2: the companion of a rational point in $[0,1]$

Definition 4.1. Let $x = p/d \in [0,1]$, where the integers $d = 1, 2, \ldots$, and p are relatively prime, in symbols, $\gcd(p,d) = 1$. Then the *companion* $\mathrm{comp}(x)$ *of* x is defined as follows: $\mathrm{comp}(0) = 1$, $\mathrm{comp}(1) = 0$, $\mathrm{comp}(1/2) = 1$; for $d = 3, 4, \ldots$, $\mathrm{comp}(p/d) = q/c$, where q/c is the neighbor (= contiguous element) of x in $\mathfrak{F}_d$ having smaller denominator than the other neighbor.

The well known unimodularity property $qd - pc = \pm 1$ of any two neighbors p/d and q/c in a Farey sequence, shows that $\mathrm{comp}(p/d)$ is unambiguously defined.

Lemma 4.2. *Adopt the above notation:*

(i) *For $d = 1, 2, 3, 4$, and $x = p/d \in [0,1]$,* $\mathrm{den}(\mathrm{comp}(x)) = 1$.

(ii) *For every $x \in \mathbb{Q} \cap [0,1]$,* $\mathrm{den}(x)$ *and* $\mathrm{den}(\mathrm{comp}(x))$ *are relatively prime.*

(iii) *For all $d \geq 3$,* $\mathrm{den}(\mathrm{comp}(p/d)) < d/2$.

Proof. We only prove (iii). Let q/c denote the companion of p/d. Let a/b be the other neighbor of p/d in $\mathfrak{F}_d$. Both segments $\mathrm{conv}(a/b, p/d)$ and $\mathrm{conv}(q/c, p/d)$ are regular. The denominators of both neighbors are smaller than d. By Lemma 3.3, a/b and q/c are neighbors in $\mathfrak{F}_{\max(b,c)}$, the segment $\mathrm{conv}(a/b, q/c)$ is regular, and $d = b + c$. (The rational p/d is known as the *Farey mediant* of a/b and q/c.) Since $d \geq 3$, then $c < b$ by definition of companion, and hence $c < d/2$. □

Conversely, we have:

Lemma 4.3. *Let d and c be integers ≥ 1. Suppose*

(i) *either $d = 1, 2, 3, 4$ and $c = 1$,*
(ii) *or $d \geq 5$ and $c < d/2$, with $\gcd(d, c) = 1$.*

Then the Farey sequence $\mathfrak{F}_d$ contains a pair of rationals p/d and q/c such that $q/c = \mathrm{comp}(p/d)$.

Proof. In case (i) the conclusion is trivial. In case (ii), let $C_0 = \mathrm{pos}[(0, 1), (1, 1)]$ be the positive hull in $\mathbb{R}^2$ of the two integer vectors $(0, 1) = u_0$ and $(1, 1) = v_0$. There is a unique sequence of cones

$$C_0 = \mathrm{pos}[u_0, v_0] \supseteq C_1 = \mathrm{pos}[u_1, v_1] \supseteq C_2 = \mathrm{pos}[u_2, v_2] \supseteq \cdots \supseteq C_t = \mathrm{pos}[u_t, v_t]$$

all containing the point (c, d) in their interior, and with the additional property that $v_{i+1} = u_i + v_i$ and $u_{i+1} \in \{u_i, v_i\}$. Each set $\{u_i, v_i\}$ is a basis of the free abelian group $\mathbb{Z}^2$. Let (u_{t1}, u_{t2}) be the coordinates of the integer vector u_t. Let similarly $v_t = (v_{t1}, v_{t2})$. Let I_i be the intersection of C_i with the line $y = 1$, and denote by J_i the projection of I_i onto the x-axis. We then obtain a decreasing sequence of regular 1-simplexes

$$J_0 = \mathrm{conv}(0, 1) \supseteq J_1 = \mathrm{conv}\left(\frac{u_{11}}{u_{12}}, \frac{v_{11}}{v_{12}}\right) \supseteq J_2 = \mathrm{conv}\left(\frac{u_{21}}{u_{22}}, \frac{v_{21}}{v_{22}}\right) \supseteq \cdots$$

By Lemma 3.3 (i)⇒(iii), the rational point c/d cannot lie in the interior of infinitely many J_i's. Thus this sequence terminates after a certain number t of steps with the segment $J_t = \mathrm{conv}(u_{t1}/u_{t2}, v_{t1}/v_{t2})$, in such a way that c/d is the Farey mediant of the vertices of J_t. We then have

$$(c, d) = (u_{t1}, u_{t2}) + (v_{t1}, v_{t2}). \tag{1}$$

Observe that $u_{t1} < u_{t2}$ and $v_{t1} < v_{t2}$. The 2×2 matrix whose columns are the vectors u_t and v_t is unimodular. Upon defining

$$q = u_{t1} \quad \text{and} \quad p = u_{t2}\,,$$

it follows that $qd - pc = \pm 1$, meaning that the 1-simplex $J = \mathrm{conv}(p/d, q/c)$ is regular. As a consequence, $\gcd(q, c) = \gcd(p, d) = 1$. By (1), $q < c$ and $p < d$, whence both p/d and q/c belong to $\mathfrak{F}_d$.

To see that q/c is a neighbor of p/d in $\mathfrak{F}_d$, arguing by way of contradiction suppose some rational $r \in \mathfrak{F}_d$ lies in the interior of J. Lemma 3.3 (i)⇒(iii) then yields $\mathrm{den}(r) \geq d + c$, against the definition of $\mathfrak{F}_d$.

To verify that q/c is the companion of p/d, let a/b be the other neighbor of p/d in $\mathfrak{F}_d$. Since the segment with vertices a/b and p/d is regular and by hypothesis $d > 1$, then $b < d$. Since $c < d$ then $\max(c, b) < d$. One more application of Lemma 3.3 now shows that q/c and a/b are contiguous elements of $\mathfrak{F}_{\max(c,b)}$, the segment $\mathrm{conv}(q/c, a/b)$ is regular, and $d = c + b$, i.e., p/d is the Farey mediant of

its neighbors. Since by hypothesis $c < d/2$ then $c < b$. Having thus shown that $q/c = \mathrm{comp}(p/d)$, the proof is complete. □

5. Rank 2: The complete classifier (G_x, c_x)

As shown by Example 3.1, when $\mathrm{rank}(G_x) = 2$ the group G_x does not suffice in general to classify $\mathrm{orb}(x)$. In Definition 5.4 we will define a numerical invariant which, jointly with the group G_x completely classifies $\mathrm{orb}(x)$. As a preliminary step we introduce the following notation and terminology:

Definition 5.1. For any fixed $n = 1, 2, 3, \ldots$ and rational affine space $\emptyset \neq F \subseteq \mathbb{R}^n$, we let $d_F = \min\{q \in \mathbb{Z} \mid q \text{ is the denominator of a rational point of } F\}$. For any $x \in \mathbb{R}^n$ we define $F_x = \bigcap\{A \subseteq \mathbb{R}^n \mid A \text{ is a rational hyperplane in } \mathbb{R}^n \text{ and } x \in A\}$.

Lemma 5.2. *For $n = 1, 2, 3, \ldots$ and $e = 0, \ldots, n$, let F be an e-dimensional rational affine space in $\mathbb{R}^n$. Let $d = d_F$.*

(i) *There are rational points $v_0, \ldots, v_e \in F$, all with denominator d, such that $\mathrm{conv}(v_0, \ldots, v_e)$ is a regular e-simplex.*
(ii) *For any rational point $y \in F$ there is $h = 1, 2, \ldots$ such that $\mathrm{den}(y) = hd$.*

Proof. (i) Starting from any rational e-simplex $R \subseteq F$ and arguing as in [13, Proposition 1], one easily obtains a regular e-simplex $\mathrm{conv}(u_0, \ldots, u_e) \subseteq R$.

For each $i = 0, \ldots, e$ the integer vector $\tilde{u}_i \in \mathbb{Z}^{n+1}$ will be more conveniently denoted $w_{0,i}$. Let F^* be the linear span in $\mathbb{R}^{n+1}$ of the primitive integer vectors $w_{0,0}, \ldots, w_{0,e}$, $F^* = \mathbb{R}w_{0,0} + \cdots + \mathbb{R}w_{0,e}$. The regularity of $\mathrm{conv}(u_0, \ldots, u_e)$ means that the set $B_0 = \{w_{0,0}, \ldots, w_{0,e}\}$ is a basis of the free abelian group $\mathbb{Z}^{n+1} \cap F^*$. If the *heights* (i.e., the last coordinates) of $w_{0,0}, \ldots, w_{0,e}$ are all equal to d we are done. Suppose not all heights of the vectors in B_0 are equal to d. We will construct a finite sequence $B_0, B_1, \ldots$ of bases of $\mathbb{Z}^{n+1} \cap F^*$, and finally obtain a basis B_{t^*} yielding the desired regular e-simplex $\mathrm{conv}(v_0, \ldots, v_e)$.

The step leading from B_0 to B_1 begins with the observation that all heights of $w_{0,0}, \ldots, w_{0,e}$ are $\geq d$, and yet it is impossible that they are all equal to the same integer $l > d$: for otherwise, a (necessarily primitive) vector in $\mathbb{Z}^{n+1} \cap F^*$ of height d, as given by the definition of d, could not arise as a linear combination of $w_{0,0}, \ldots, w_{0,e}$ with integer coefficients, and B_0 would not be a basis of $\mathbb{Z}^{n+1} \cap F^*$. Since not all vectors of B_0 have the same height, we pick in B_0 a vector $w_{0,i}$ of largest height, a vector $w_{0,j}$ of smallest height, and replace $w_{0,i}$ by the new integer vector $w_{0,i} - w_{0,j}$ leaving the rest unchanged. We get a new basis $B_1 = \{w_{1,0}, \ldots, w_{1,e}\}$ of $\mathbb{Z}^{n+1} \cap F^*$. The height of each element of B_1 is still $\geq d$. The sum of the heights of the elements of B_1 is strictly smaller than the sum of the heights of the elements of B_0. The positive hull of these vectors yields an $(e+1)$-dimensional regular cone C_1, from which we get a regular e-simplex $\mathrm{conv}(w_{1,0}^{\downarrow}, \ldots, w_{1,e}^{\downarrow}) \subseteq \mathbb{R}^n$ by letting $w_{1,i}^{\downarrow}$ be the only rational point in $\mathbb{R}^n$ such that $w_{1,i}$ is the homogeneous correspondent of $w_{1,i}^{\downarrow}$.

Proceeding inductively, suppose we have a basis $B_t = \{w_{t,0}, \ldots, w_{t,e}\}$ of $\mathbb{Z}^{n+1} \cap F^*$, whose positive hull is the cone $C_t \subseteq \mathbb{R}^{n+1}$, and its associated regular e-simplex is $\mathrm{conv}(w_{t,0}^{\downarrow}, \ldots, w_{t,e}^{\downarrow}) \subseteq \mathbb{R}^n$. If the heights of all the vectors of B_t are equal to d we are done. Otherwise, as above, we are allowed to pick in B_t a top vector $w_{t,i}$, a bottom vector $w_{t,j}$, and replace $w_{t,i}$ by $w_{t,i} - w_{t,j}$ thus obtaining a new basis $B_{t+1} = \{w_{t+1,0}, \ldots, w_{t+1,e}\}$. From B_{t+1} we get a new regular cone $C_{t+1} \subseteq \mathbb{R}^{n+1}$

and a new regular e-simplex $\mathrm{conv}(w^{\downarrow}_{t+1,0},\dots,w^{\downarrow}_{t+1,e})\subseteq\mathbb{R}^n$ by the same procedure of the first step. The height of each element of B_{t+1} is still $\geq d$. Since the sum of the heights of the elements of B_{t+1} is strictly smaller than the sum of the heights of the elements of B_t, this sequence terminates with a basis B_{t^*} of $\mathbb{Z}^{n+1}\cap F^*$ whose elements all have the same height d. From B_{t^*} we obtain the regular cone C_{t^*} and the desired regular e-simplex $\mathrm{conv}(v_0,\dots,v_e)$.

(ii) is a trivial consequence of (i): for, the primitive vector $\tilde{y}$ is a linear combination of the $\tilde{v}_i$ with integer coefficients, and $1\leq\mathrm{den}(y)\in\mathbb{Z}$. □

The following characterization of d_{F_x} will find repeated use in the sequel:

Lemma 5.3. *With reference to Definition 5.1, for every $x\in\mathbb{R}^2$ with $\mathrm{rank}(G_x)=2$ let us write for short d instead of d_{F_x}. Let us display F_x as the rational line $\{(x_1,x_2)\in\mathbb{R}^2\mid a_1x_1+a_2x_2+a_3=0\}$, for relatively prime integers a_1,a_2,a_3. Noting that the primitive integer vector $(a_1,a_2,a_3)\in\mathbb{Z}^3$ is uniquely determined up to scalar multiplication by ±1 we then have:*

(i) $d=\gcd(a_1,a_2)$.

(ii) *d is the denominator of the smallest nonzero positive rational in G_x.*

Proof. (i) Let us use the abbreviations $D=\gcd(a_1,a_2)$, $a'_1=a_1/D$, $a'_2=a_2/D$. Then $\gcd(D,a_3)=\gcd(a'_1,a'_2)=1$ and for suitable integers α,β we can write

$$\alpha a'_1+\beta a'_2=-1,\ \ \gcd(\alpha,\beta)=1.$$

Thus $a_1(\alpha a_3/D)+a_2(\beta a_3/D)=-a_3$, and the rational point $(\alpha a_3/D,\beta a_3/D)$ belongs to F_x. From $\gcd(D,a_3)=\gcd(\alpha,\beta)=1$ it follows that $\mathrm{den}(\alpha a_3/D,\beta a_3/D)=D$.

To conclude the proof that $d=D$, we *claim* that no rational point $z\in\mathbb{R}^2$ with $\mathrm{den}(z)<D$ lies in F_x. For otherwise, for suitable integers m,n,δ, let $z=(m/\delta,n/\delta)$ be a counterexample (absurdum hypothesis). From $a_1m/\delta+a_2n/\delta=-a_3$ we get $a_1m+a_2n=-a_3\delta$. On the other hand, $a_1m+a_2n=kD$ for some integer k. It follows that $a_3\delta=kD$. Since $\delta<D$ some prime divisor of D must also be a divisor of a_3, which contradicts $\gcd(D,a_3)=1$. Our claim is settled, and hence $d=D=\gcd(a_1,a_2)$, as required to complete the proof of (i).

(ii) Let $H=\{\tilde{y}\in\mathbb{Z}^3\mid y\in F_x\cap\mathbb{Q}\}$ be the set of homogeneous correspondents of all rational points of F_x. Then every point $z\in F_x$ of denominator d corresponds to an integer point $\tilde{z}\in H$ of height d. One then sees that points of denominator d lying on the line F_x are equidistant. Each segment in F_x joining any two consecutive such points is regular: this follows from Lemma 3.3(iii)⇒(i) in combination with Lemma 5.2. The point x lies in the convex hull of precisely two points $a,b\in F_x$ of denominator d.

Pick a rational point $s\notin F_x$ such that the triangle $T=\mathrm{conv}(a,b,s)$ is regular. Let $c=\mathrm{den}(s)$. The regularity of T is to the effect that $\gcd(d,c)=1$. Then a routine variant of the argument in Lemma 4.3 yields rationals q/c and p/d (written in their lowest terms) which are neighbors in the Farey sequence $\mathfrak{F}_{\max(c,d)}$. The triangle $U=\mathrm{conv}((1/d,0),(0,p/d),(0,q/c))$ is regular. By Lemma 2.2 there is a map $\gamma\in\mathcal{G}_2$ such that $\gamma(T)=U$. In particular $\gamma(x)=\xi=(\xi_1,\xi_2)$ for suitable irrationals ξ_1 and ξ_2. It follows that $G_x=G_\xi$ is generated by $\xi_1,\xi_2,1$. The point ξ lies on the line F_ξ. Thus $\xi_2=p(1/d-\xi_1)$. We see that $p/d\in G_x$. Since $1\in G_x$, from $\gcd(p,d)=1$ we obtain $1/d\in G_x$.

To conclude the proof, by way of contradiction let u/v be a rational in G_x with

$$0 < u/v < 1/d. \tag{2}$$

From $\gcd(u,v) = 1$ and $1 \in G_x$ we obtain $1/v \in G_x$, whence $1/\operatorname{lcm}(d,v) \in G_x$, where lcm denotes least common multiple. Upon writing $\operatorname{lcm}(v,d) = kd$ for some integer k, since $\operatorname{rank}(G_x) = 2$ then $\{1/d, \xi_1\}$ is a *free* generating set of G_x, and there are integers m, n such that $m\xi_1 + n/d = 1/(kd)$, whence $nk = 1$, $n = k = 1$, and $v = d$. This contradicts (2). □

We are now ready to introduce the numerical invariant which, in combination with $G(x)$, yields a complete invariant for $\operatorname{orb}(x)$.

Definition 5.4. For any $n = 1, 2, \ldots$ and e-dimensional rational affine space F in $\mathbb{R}^n$ $(e = 0, 1, \ldots, n)$, we define c_F as the smallest possible denominator of a vertex of a regular n-simplex $\operatorname{conv}(v_0, \ldots, v_n) \subseteq \mathbb{R}^n$ with $v_0, \ldots, v_e \in F$.

For every $x \in \mathbb{R}^n$ we write c_x as an abbreviation of c_{F_x}.

The two points a, b of Example 3.1 have $c_a = 1$ and $c_b = 2$.

Lemma 5.5. *Let $n = 2, 3, \ldots$ and $x \in \mathbb{R}^n$.*

(i) *If $\operatorname{rank}(G_x) = n + 1$, i.e., if F_x is n-dimensional, then $c_x = 1$.*
(ii) *If $\operatorname{rank}(G_x) = 1$, i.e., if F_x is 0-dimensional, then $c_x = 1$.*
(iii) *Suppose F_x is e-dimensional, with $0 < e < n$, meaning that $\operatorname{rank}(G_x) \notin \{1, n+1\}$. Write for short d instead of d_{F_x}. If $d \in \{1, 2\}$ then $c_x = 1$. If $d \geq 3$ then $c_x < d/2$ and $\gcd(d, c_x) = 1$.*

Proof. (i) The hypothesis means that x is totally irrational. Then $F_x = \mathbb{R}^n$. It is easy to construct a regular simplex T in $\mathbb{R}^n$ such that the denominator of each vertex of T is equal to 1. By definition, $c_x = 1$.

(ii) By Remark 2.4.

(iii) Let $V = \operatorname{conv}(v_0, \ldots, v_n) \subseteq \mathbb{R}^n$ be a regular n-simplex with $v_0, \ldots, v_e \in F_x$, and with a vertex v_i satisfying $\operatorname{den}(v_i) = c_x$.

Claim 1. There is a regular n-simplex $W = \operatorname{conv}(w_0, \ldots, w_n) \subseteq \mathbb{R}^n$ whose vertices $w_0, \ldots, w_e$ lie in F_x, and with $\operatorname{den}(w_n) = c_x$.

As a matter of fact, if $\operatorname{den}(v_j) = c_x$ for some $j = e+1, \ldots, n$, then without loss of generality $j = n$, and we are done. Otherwise, for all $j = e+1, \ldots, n$ we have the inequality $c_x < \operatorname{den}(v_j)$, and we can write $\operatorname{den}(v_0) = c_x$ without loss of generality. Passing to homogeneous correspondents in $\mathbb{R}^{n+1}$, for some integer $k > 0$ the height (= last coordinate) l of the vector $r_n = \tilde{v}_n - k\tilde{v}_0 \in \mathbb{R}^{n+1}$ will satisfy $1 \leq l \leq$ the height of $\tilde{v}_0$. Thus by definition of c_x, both r_n and $\tilde{v}_0$ have the same height c_x. Let $z_n \in \mathbb{R}^n$ be the rational point whose homogeneous correspondent $\tilde{z}_n$ equals r_n. Then $\operatorname{den}(z_n) = c_x =$ height of r_n. The regular n-simplex $W = \operatorname{conv}(v_0, \ldots, v_{n-1}, z_n)$ satisfies the conditions of our claim.

An application of Lemmas 5.2 and 3.2 now yields

Claim 2. There is a regular n-simplex $Y = \operatorname{conv}(y_0, \ldots, y_n) \subseteq \mathbb{R}^n$ with $y_0, \ldots, y_e \in F_x$, satisfying the identities $\operatorname{den}(y_n) = c_x$ and $\operatorname{den}(y_0) = \cdots = \operatorname{den}(y_e) = d$.

We conclude the proof arguing by cases:

If $d = 1$ we are done, because $1 \leq c_x \leq d$.

If $d = 2$ we *claim* that $c_x = 1$. If not (absurdum hypothesis), $c_x = 2$ and not all of $\operatorname{den}(y_{e+1}), \ldots, \operatorname{den}(y_n)$ can be even, because the regularity of the n-simplex

$Y = \mathrm{conv}(y_0, \ldots, y_n)$ entails the unimodularity of the matrix whose columns are $\tilde{y}_0, \ldots, \tilde{y}_n$. Thus for some $i = e+1, \ldots, n$, $\mathrm{den}(y_i)$ is odd and ≥ 3. For some integer $k \geq 1$, the height of the integer vector $l_i = \tilde{y}_i - k\tilde{y}_0 \in \mathbb{R}^{n+1}$ is equal to 1. Let the rational point $u_i \in \mathbb{R}^n$ be defined by $\tilde{u}_i = l_i$. Replacing in Y the vertex y_i by u_i, the resulting n-simplex $\mathrm{conv}(y_0, \ldots, y_{i-1}, u_i, y_{i+1} \ldots, y_n)$ is regular. Since $\mathrm{den}(u_i) = 1$ then $c_x = 1$, a contradiction with our absurdum hypothesis $c_x = 2$. We conclude that $c_x = 1$, as required to settle our claim.

If $d \geq 3$, for every $l = e+1, \ldots, n-1$ such that $\mathrm{den}(y_l) > c_x$ there is an integer $k_l \geq 1$ such that the height h of the vector $s_l = \tilde{y}_l - k_l\tilde{y}_n \in \mathbb{Z}^{n+1}$ satisfies $1 \leq h \leq c_x$, whence $h = c_x$. Letting the rational point u_l be defined by $\tilde{u}_l = s_l$, replace vertex $y_l \in Y$ by u_l. We then obtain a new regular n-simplex with at least two vertices not in F_x, both having denominator equal to c_x. This process terminates with a regular n-simplex $Z = \mathrm{conv}(y_0, \ldots, y_e, u_{e+1}, \ldots, u_{n-1}, y_n)$ whose first $e+1$ vertices lie in F_x and have denominator d, and whose remaining vertices have denominator c_x and lie outside F_x. The regularity of Z entails that c_x and d are relatively prime. Since $d > 1$ then $c_x < d$. Since c_x is minimal, then $c_x \leq d/2$. Actually, $c_x < d/2$. For otherwise, d is even, $d \geq 4$, and $\gcd(c_x, d) = c_x \geq 2$, contradicting the regularity of Z. □

6. Rank 2: the classification theorem

Our classification of $\mathcal{G}_2$-orbits is completed as follows:

Theorem 6.1. *If* $x \in \mathbb{R}^2$ *and* $\mathrm{rank}(G_x) = 2$ *then* $\mathrm{orb}(x) = \{x' \in \mathbb{R}^2 \mid (G_x, c_x) = (G_{x'}, c_{x'})\}$.

Proof. The rational affine subspace $F_x \subseteq \mathbb{R}^2$ is the only rational line in $\mathbb{R}^2$ passing through x. We first prove $\mathrm{orb}(x) \subseteq \{x' \in \mathbb{R}^2 \mid G_x = G_{x'} \text{ and } c_x = c_{x'}\}$. If x and x' satisfy $\delta(x) = x'$ for some $\delta \in \mathcal{G}_2$, then we immediately see that $G_x = G_{x'}$. Further, the map γ sends F_x one-one onto $F_{x'}$. Trivially, γ preserves denominators of rational points. Lemma 3.3(i)⇔(iii) shows that γ also induces a one-one correspondence between regular simplexes. Thus $c_x = c_{x'}$.

For the converse inclusion $\mathrm{orb}(x) \supseteq \{x' \in \mathbb{R}^2 \mid G_x = G_{x'} \text{ and } c_x = c_{x'}\}$, let us suppose $G_x = G_{x'}$ and $c_x = c_{x'}$, looking for an affine transformation $\delta \in \mathcal{G}_2$ such that $\delta(x) = x'$. For suitable integers a_1, a_2, a_3 with $\gcd(a_1, a_2, a_3) = 1$, and a_1 and a_2 not both zero, we can write $F_x = \{(y_1, y_2) \in \mathbb{R}^2 \mid a_1y_1 + a_2y_2 + a_3 = 0\}$. By Lemma 5.3, d_{F_x} coincides with the denominator of the smallest positive nonzero rational point of G_x. Let us write d instead of d_{F_x}. The points of denominator d lying on the line F_x are at equal distance. Each segment in F_x joining any two consecutive such points is regular by Lemmas 3.3(iii)⇒(i) and 5.2(ii). The point x lies in the convex hull of precisely two points $a, b \in F_x$ of denominator d.

The same construction in the proof of Lemma 5.5(iii) yields a rational point $s \notin F_x$ such that the triangle $T = \mathrm{conv}(a, b, s)$ is regular, and $\mathrm{den}(s) = c_x$. Let us write c instead of c_x.

The denominators of a, b, s are equal to d, d, c respectively. As a consequence of the regularity of T, $\gcd(c, d) = 1$. By Lemmas 5.5 the integers c and d satisfy the hypotheses of Lemma 4.3, whence we obtain integers $p < d$ and $q < c$ such that $q/c = \mathrm{comp}(p/d)$ and

$$qd - pc = \pm 1. \tag{3}$$

Let the triangle $\bar{T} = \mathrm{conv}(\bar{a}, \bar{b}, \bar{s})$ be defined by $\bar{a} = (1/d, 0), \bar{b} = (0, p/d), \bar{s} = (0, q/c)$. Then $\bar{T}$ is regular and the denominators of its vertices are respectively equal to d, d, c. By Lemma 2.2 some $\alpha \in \mathcal{G}_2$ maps a, b, s to $\bar{a}, \bar{b}, \bar{s}$. For suitable irrational numbers ξ_1, ξ_2 we can write $\alpha(x) = \bar{x} = (\xi_1, \xi_2) \in \mathrm{conv}(\bar{a}, \bar{b})$. Evidently, $G_x = G_{\bar{x}}$, the latter being the subgroup of $\mathbb{R}$ generated by $\xi_1, \xi_2, 1$. Since $\bar{x} \in F_{\bar{x}} = \alpha(F_x)$ then

$$\xi_2 = p(1/d - \xi_1). \tag{4}$$

Thus the subgroup H of $\mathbb{R}$ generated by ξ_1 and $1/d$ contains $G_{\bar{x}}$ as a subgroup, $G_{\bar{x}} \subseteq H$. Conversely, $H \subseteq G_{\bar{x}}$. As a matter of fact, from $\gcd(p, d) = 1$ it follows that $lp + md = 1$ for suitable integers l, m. Thus $l \cdot p/d + m \cdot 1 = 1/d$, and hence $1/d \in G_{\bar{x}}$, because $p/d \in G_{\bar{x}}$, by (4). We have shown $G_x = G_{\bar{x}} = H = \mathbb{Z}\xi_1 + \mathbb{Z}\frac{1}{d}$. From the hypothesis $\mathrm{rank}(G_x) = 2$ it follows that

$$G_x \text{ is freely generated by } \xi_1 \text{ and } 1/d. \tag{5}$$

We now consider the point x' and its associated group $G_{x'} = G_x$. By Lemma 5.3 the integer $d' = d_{F_{x'}}$ coincides with the denominator of the smallest positive nonzero rational point of $G_{x'}$. Since $G_{x'} = G_x$ then $d' = d$. By hypothesis, $c_x = c_{x'} = c$. The point x' lies in the convex hull of precisely two points $a', b' \in F_{x'}$ of denominator d. As above, there is a rational point $s' \notin F_{x'}$ such that the triangle $T' = \mathrm{conv}(a', b', s')$ is regular, and $\mathrm{den}(s') = c$. Since the denominators of the vertices a', b', s' are equal to d, d, c, some $\beta \in \mathcal{G}_2$ will map T' onto *the same* triangle $\alpha(T) = \mathrm{conv}(\bar{a}, \bar{b}, \bar{s})$. In particular, $\beta(x') = \bar{x}'$ for some point $\bar{x}' = (\xi_1', \xi_2') \in \mathrm{conv}(\bar{a}, \bar{b})$. It follows that

$$G_{x'} = G_x \text{ is freely generated by } \xi_1' \text{ and } 1/d. \tag{6}$$

Recalling (5), some 2×2 integer unimodular matrix U maps the column vector $(\xi_1, 1/d)$ to the column vector $(\xi_1', 1/d)$. Since ξ_1 is irrational, the bottom row of U has the form $(0, 1)$. So, for some $h = \pm 1$,

$$\begin{pmatrix} h & k \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \xi_1 \\ 1/d \end{pmatrix} = \begin{pmatrix} \xi_1' \\ 1/d \end{pmatrix} \tag{7}$$

To get the desired map $\delta \in \mathcal{G}_2$ sending x to x' we have only to exhibit a map $\gamma \in \mathcal{G}_2$ with $\gamma(\bar{x}) = \bar{x}'$.

Case 1: $h = 1$. Then $k = 0$, because $0 < \xi_1' < 1/d$, and $0 < \xi_1 < 1/d$. It follows that $\bar{x} = \bar{x}'$ and the identity map on $\mathbb{R}^2$ provides the required $\gamma \in \mathcal{G}_2$.

Case 2: $h = -1$. Then from $\xi_1' = -\xi_1 + k/d$ we immediately obtain $k = 1$, because $0 < \xi_1' < 1/d$, and $0 < \xi_1 < 1/d$. From (7) have the identities

$$\xi_1' = -\xi_1 + 1/d, \quad \xi_2' = p\xi_1. \tag{8}$$

Since ξ_2' lies in F_x,

$$\xi_2' = p(1/d - \xi_1'). \tag{9}$$

To conclude the proof, recalling (3), let $u = cp - dq = \pm 1$. Let the integer matrix M be given by

$$M = \begin{pmatrix} dq/u & c/u & -q/u \\ p - dpq/u & -dq/u & pq/u \\ 0 & 0 & 1 \end{pmatrix}$$

Then M is unimodular and maps $(\xi_1, \xi_2, 1)$ to $(\xi_1', \xi_2', 1)$. To see this, one verifies the identity

$$M \begin{pmatrix} \xi_1 \\ p(1/d - \xi_1) \\ 1 \end{pmatrix} = \begin{pmatrix} -\xi_1 + 1/d \\ p\xi_1 \\ 1 \end{pmatrix}$$

using (4)-(6) and (8)-(9). From M we immediately obtain the desired $\gamma \in \mathcal{G}_2$ satisfying $\gamma(\bar{x}) = \bar{x}'$. □

The following result states that the pair (G_x, c_x) is a complete invariant for the orbit of $x \in \mathbb{R}^2$:

Corollary 6.2. *Two points $x, x' \in \mathbb{R}^2$ have the same orbit iff $G_x = G_{x'}$ and $c_x = c_{x'}$.*

Proof. The case $\mathrm{rank}(G_x) = 2$ is taken care of by the foregoing result. The cases $\mathrm{rank}(G_x) \in \{1, 3\}$ are dealt with by Lemma 5.5(i)-(ii). Here c_x is automatically equal to 1, and its role in the classification of $\mathrm{orb}(x)$ is immaterial. □

Corollary 6.3. *For $x = (x_1, x_2)$ ranging over all points in $\mathbb{R}^2$ let the subgroup G_x of $\mathbb{R}$ be defined by $G_x = \mathbb{Z} + \mathbb{Z}x_1 + \mathbb{Z}x_2$. Let c_x be as in Definition 5.4. Then the possible classifiers (G_x, c_x) of $\mathcal{G}_2$-orbits are as follows:*

(i) *If $\mathrm{rank}\, G_x \in \{1, 3\}$ then automatically $c_x = 1$.*
(ii) *If $\mathrm{rank}\, G_x = 2$, let d be the denominator of the smallest nonzero positive rational element of G_x. If $d \leq 4$ then $c_x = 1$. If $d = 5, 6, \ldots$ then any integer c relatively prime with d and such that $0 < c < d/2$ may occur as c_x. Thus precisely $\max(1, \phi(d)/2)$ distinct orbits are classified by the same group G_x.*
(iii) *No other pair (G_x, c_x) may occur.*

Proof. From Lemmas 4.2, 4.3, 5.5 and Theorem 6.1. □

The complete classification of $\mathcal{G}_1$-orbits is now easily obtained as follows:

Corollary 6.4. *For x ranging over all points in $\mathbb{R}$ let the subgroup G_x of $\mathbb{R}$ be defined by $G_x = \mathbb{Z} + \mathbb{Z}x$. Let c_x be as in Definition 5.4. Then the possible classifiers (G_x, c_x) of $\mathcal{G}_1$-orbits are as follows:*

(i) *In case $\mathrm{rank}(G_x) = 2$ then $c_x = 1$. For all $y \in \mathbb{R}$, $\mathrm{orb}(x) = \mathrm{orb}(y)$ iff $G_x = G_y$.*

In case $\mathrm{rank}(G_x) = 1$, letting d denote the denominator of the smallest nonzero positive rational element of G_x,

(ii) *If $d \leq 4$, then $c_x = 1$. For all $y \in \mathbb{R}$, $\mathrm{orb}(x) = \mathrm{orb}(y)$ iff $G_x = G_y$.*
(iii) *If $d > 4$, precisely $\max(1, \phi(d)/2)$ distinct orbits are classified by the same group G_x. These orbits are in one-one correspondence with the totality of pairs (G_x, c) with $c \in \mathbb{Z}$, $1 \leq c < d/2$ and $\gcd(c, d) = 1$.*

Concerning the topological properties of $\mathrm{orb}(x)$, considerations from ergodic theory and dynamics of flows in the homogeneous space $SL(n,\mathbb{R})/SL(n,\mathbb{Z})$ lead to the well known result [3, 5.2] which, in the framework of the present paper, can be reformulated as follows: *for each* $n = 2, 3, \ldots$ *and* $x \in \mathbb{R}^n$, $\mathrm{orb}(x)$ *is dense iff* $\mathrm{rank}(G_x) \neq 1$.

7. Applications

Applications to lattice-ordered abelian groups with order unit. In this section we assume familiarity with lattice-ordered abelian groups. We refer to [1] for background. An *ℓ-group* (which is short for "lattice-ordered abelian group") is an abelian group G equipped with a translation invariant lattice order. If G also has a distinguished (strong) *order unit*, i.e., an element u whose positive integer multiples eventually dominate any element of G, then (G,u) is known as a *unital ℓ-group*. Morphisms in the category of unital ℓ-groups are *unital ℓ-homomorphisms*, i.e., group homomorphisms that also preserve the lattice structure and the distinguished order unit. We write $(G,u) \cong (H,v)$ when (G,u) and (H,v) are unitally ℓ-isomorphic.

An *ideal* ("ℓ-ideal" in [1]) $\mathfrak{i}$ of (G,u) is the kernel of a unital ℓ-homomorphism $\eta\colon (G,u) \to (H,v)$. In case $\eta(G,u)$ is totally ordered, $\mathfrak{i}$ is said to be *prime* ("irréductible" in [1]). Hölder's theorem, [1, 2.6] is to the effect that for every maximal ideal $\mathfrak{m}$ of (G,u) the quotient $(G,u)/\mathfrak{m}$ is uniquely embeddable into a unital ℓ-subgroup of the additive group $\mathbb{R}$ equipped with the usual order and with the element 1 as the order unit. The set $\mu(G)$ of maximal ideals of (G,u) is a subset of the set of prime ideals. The hull-kernel topology makes $\mu(G,u)$ a compact Hausdorff space, [1, 13.2.6]. We then say that $\mu(G,u)$ is the *maximal spectral space* of (G,u).

Since the category of ℓ-groups is equivalent to an equational class, [11, 3.9], one can naturally introduce the free unital ℓ-group $\mathcal{F}_n$ over n free generators. Every finitely generated unital ℓ-group is (isomorphic to) the quotient $\mathcal{F}_n/\mathfrak{i}$ of some $\mathcal{F}_n$ by an ideal $\mathfrak{i}$. The maximal spectral space of $\mathcal{F}_n$ is homeomorphic to the unit cube $[0,1]^n$ with the product topology, so it makes sense to say that a maximal ideal $\mathfrak{m}$ of $\mathcal{F}_n$ "lies in the interior" of $\mu(\mathcal{F}_n)$. See [13, Proposition 3(i)] for details.

As a particular case of [1, 10.5.3], for every maximal ideal $\mathfrak{m}$ of (G,u) by the *germinal* ideal $\mathfrak{o}(\mathfrak{m})$, we mean the intersection of all prime ideals $\mathfrak{p}$ of (G,u) contained in $\mathfrak{m}$.

Theorem 7.1. *Let $\mathcal{F}_2$ be the free unital ℓ-group on two generators. Let $\mathfrak{m}$ be a maximal ideal of $\mathcal{F}_2$ lying in the interior of $\mu(\mathcal{F}_2)$. Write $G_\mathfrak{m}$ for the underlying group of the unital ℓ-group $\mathcal{F}_2/\mathfrak{m}$.*

(i) *$G_\mathfrak{m}$ is a subgroup of the additive group $\mathbb{R}$ of rank ≤ 3 containing 1.*

(ii) *If $\mathrm{rank}(G_\mathfrak{m}) = 3$ then for every maximal ℓ-ideal $\mathfrak{n}$,*

$$\mathcal{F}_2/\mathfrak{m} \cong \mathcal{F}_2/\mathfrak{n} \Leftrightarrow \mathcal{F}_2/\mathfrak{o}(\mathfrak{m}) \cong \mathcal{F}_2/\mathfrak{o}(\mathfrak{n}).$$

If $\mathrm{rank}(G_\mathfrak{m})$ equals 1 then for every maximal ℓ-ideal $\mathfrak{n}$ lying in the interior of $\mu(\mathcal{F}_2)$,

$$\mathcal{F}_2/\mathfrak{m} \cong \mathcal{F}_2/\mathfrak{n} \Leftrightarrow \mathcal{F}_2/\mathfrak{o}(\mathfrak{m}) \cong \mathcal{F}_2/\mathfrak{o}(\mathfrak{n}).$$

(iii) *If $\mathrm{rank}(G_\mathfrak{m}) = 2$, let d be the denominator of the smallest positive nonzero rational in $G_\mathfrak{m}$. Then the set of isomorphism classes of unital ℓ-groups of*

the form $\mathcal{F}_2/\mathfrak{o}(\mathfrak{n})$, *for* $\mathfrak{n}$ *a maximal ideal in the interior of* $\mu(\mathcal{F}_2)$ *such that* $\mathcal{F}_2/\mathfrak{n} \cong \mathcal{F}_2/\mathfrak{m}$, *contains exactly* $\max\left(1, \frac{\phi(d)}{2}\right)$ *elements.*

Proof. By [13, Proposition 2], $\mathcal{F}_2$ has a concrete representation as the unital ℓ-group of all piecewise linear continuous functions $f\colon [0,1]^2 \to \mathbb{R}$, where each piece of f is a linear (affine) polynomial with integer coefficients. The constant function 1 is the distinguished order unit of $\mathcal{F}_2$. The operations of $\mathcal{F}_2$ are pointwise addition, subtraction, max, and min. Via the Yosida correspondences $\mathfrak{n} \mapsto x_{\mathfrak{n}}$ and $x \mapsto \mathfrak{m}_x$ of [13, Proposition 3(i)] we further identify the maximal spectral space $\mu(\mathcal{F}_2)$ with $[0,1]^2$, in such a way that the maximal ideal $\mathfrak{m}_x$ is the set of all functions vanishing at $x_{\mathfrak{m}}$, and the point $x_{\mathfrak{n}}$ is the intersection of the zerosets of all functions in $\mathfrak{n}$. For every $\mathfrak{p} \in \mu(\mathcal{F}_2)$, the quotient map $f \in \mathcal{F}_2 \mapsto f/\mathfrak{p}$ amounts to evaluation at $x_{\mathfrak{p}}$,

$$f/\mathfrak{p} = f(x_{\mathfrak{p}}).$$

Then the germinal ideal $\mathfrak{o}(\mathfrak{n}_x)$ is the set of all functions vanishing on an open neighbor of $x_{\mathfrak{p}}$.

(i) By Hölder's theorem, $G_{\mathfrak{m}}$ is the subgroup of $\mathbb{R}$ generated by the unit 1 together with the coordinates x_1, x_2 of x. So $G_{\mathfrak{m}} = G_x$ and $\mathrm{rank}(G_{\mathfrak{m}}) \in \{1,2,3\}$. The desired conclusion now follows from [13, Proposition 3(ii)-(iii)].

(ii) In case $\mathrm{rank}(G_{\mathfrak{m}}) = 3$, the point $x_{\mathfrak{m}}$ is totally irrational and thus automatically lies in the interior of $\mu(\mathcal{F}_2)$. Each function $f \in \mathcal{F}_2$ vanishing at $x_{\mathfrak{m}}$ also vanishes on an open neighborhood of $x_{\mathfrak{m}}$. Therefore, the maximal quotient $\mathcal{F}_2/\mathfrak{m}$ coincides with the germinal quotient $\mathcal{F}_2/\mathfrak{o}(\mathfrak{m})$, and we are done.

In case $\mathrm{rank}(G_{\mathfrak{m}}) = 1$, one first proves the following claim: *There is an isomorphism* $\iota\colon \mathcal{F}_2/\mathfrak{o}(\mathfrak{m}) \cong \mathcal{F}_2/\mathfrak{o}(\mathfrak{n})$ *iff (*$\mathfrak{n}$ *lies in the interior of* $\mu(\mathcal{F}_2)$ *and) there exist rational triangles* T *and* U *with* $x_{\mathfrak{m}} \in \mathrm{int}(T)$ *and* $y_{\mathfrak{n}} \in \mathrm{int}(U)$ *and a map* $\gamma \in \mathcal{G}_2$ *satisfying* $\gamma(x_{\mathfrak{m}}) = x_{\mathfrak{n}}$ *and* $\gamma(T) = U$. By Proposition 2.3, a necessary and sufficient condition for such γ to exist is that $G_{\mathfrak{m}} = G_{\mathfrak{n}}$. By Hölder's theorem, this is in turn equivalent to saying that the unital ℓ-groups $\mathcal{F}_2/\mathfrak{m}$ and $\mathcal{F}_2/\mathfrak{n}$ are unitally ℓ-isomorphic.

(iii) From Lemma 5.5(iii) it follows that, whenever $\mathrm{rank}(G_x) = 2$, the number of orbits classified by the same group $G_{\mathfrak{m}}$ is $\max(1, \phi(d)/2)$, and by Theorem 6.1 these orbits are in one-one correspondence with the possible values of the parameter $c_{x_{\mathfrak{m}}}$ of Definition 5.4. □

Applications to the Farey-Stern-Brocot AF C*-algebra. A *(unital) AF C*-algebra* $\mathfrak{A}$ is the norm closure of an ascending sequence of finite-dimensional C*-algebras, all with the same unit. In this section we require acquaintance with AF C*-algebras and Elliott classification [5, 7],

$$\mathfrak{A} \mapsto (K_0(\mathfrak{A}), [1_{\mathfrak{A}}]),$$

of unital AF C^*-algebras in terms of countable unital dimension groups. Here, K_0 is a suitable order-theoretic expansion of Grothendieck's functor, and $[1_{\mathfrak{A}}]$ is the K_0-image of the unit of $\mathfrak{A}$.

The Murray-von Neumann order of projections in $\mathfrak{A}$ is a lattice iff $K_0(\mathfrak{A})$ is an ℓ-group. Combining Corollary 6.4 with Elliott's classification, in Theorem 7.2 we will describe the mutual relationships between the maximal and the germinal ideals of the Farey-Stern-Brocot C^*-algebra $\mathfrak{M}_1$ introduced by the present author

in [12] and recently rediscovered by Boca in [2] (also see [4] and [14]). The Bratteli diagram of $\mathfrak{M}_1$ is as follows:



The two top (depth 0) vertices are labeled 1. The label of any vertex v at depth $d = 1, 2, \ldots$, is the sum of the labels of the vertices at depth $d-1$ connected to v by an edge.

As proved in [12], an equivalent redefinition of $\mathfrak{M}_1$ is as follows:

$$(K_0(\mathfrak{M}_1), [1_{\mathfrak{M}_1}]) = \mathcal{F}_1.$$

Now, up to unital ℓ-isomorphism, $\mathcal{F}_1$ is the unital ℓ-group of continuous piecewise linear functions $f\colon [0,1] \to \mathbb{R}$, where each piece of f is a linear polynomial with integer coefficients, [13, Proposition 2]. The constant function 1 is the distinguished order unit of $\mathcal{F}_1$. As a particular case of a general well known fact [14, §3], the transformation

$$\mathfrak{P} \mapsto K_0(\mathfrak{P}) \tag{10}$$

induces an order preserving one-one map of the ideals of $\mathfrak{M}_1$ onto the ideals of $\mathcal{F}_1$, in such a way that

$$K_0\left(\frac{(\mathfrak{M}_1, [1_{\mathfrak{M}_1}])}{\mathfrak{P}}\right) = \frac{K_0(\mathfrak{M}_1, [1_{\mathfrak{M}_1}])}{K_0(\mathfrak{P})}. \tag{11}$$

In particular, K_0 sends primitive (resp., maximal) ideals of $\mathfrak{M}_1$ one-one onto prime (resp., maximal) ideals of $\mathcal{F}_1$. For every maximal ideal $\mathfrak{I}$ of of $\mathfrak{M}_1$, the *germinal ideal* $\mathfrak{O}(\mathfrak{I})$ of $\mathfrak{M}_1$ at $\mathfrak{I}$ is the intersection of all primitive ideals of $\mathfrak{M}_1$ contained in $\mathfrak{I}$. The K_0-image of $\mathfrak{O}(\mathfrak{I})$ is the germinal ideal $\mathfrak{o}(K_0(\mathfrak{I}))$.

For each irrational θ, the AF C*-algebra $\mathfrak{F}_\theta$ is defined by

$$K_0(\mathfrak{F}_\theta, [1_{\mathfrak{F}_\theta}]) = (\mathbb{Z} + \mathbb{Z}\theta, 1) \tag{12}$$

i.e., the subgroup of the additive group $\mathbb{R}$ generated by 1 and θ, equipped with the usual order, and with the distinguished order unit 1. See [5, p. 34, and p.65] for further information.

As proved in [14, 3.4], the maximal ideal space of $\mathfrak{M}_1$ equipped with the hull-kernel topology is canonically homeomorphic to the unit real interval $[0,1]$.

Theorem 7.2. *Suppose $\mathfrak{I}$ is a maximal ideal of $\mathfrak{M}_1$.*

(i) *If $\mathfrak{M}_1/\mathfrak{I} \cong \mathfrak{F}_\theta$ for some irrational $\theta \in [0,1]$, then $\mathfrak{M}_1/\mathfrak{I} \cong \mathfrak{M}_1/\mathfrak{O}(\mathfrak{I})$.*

(ii) *If* $\mathfrak{M}_1/\mathfrak{I} \ncong \mathfrak{F}_\theta$ *for every irrational* $\theta \in [0,1]$, *then* $\mathfrak{M}_1/\mathfrak{I}$ *is isomorphic to the* C^**-algebra of linear operators on* d*-dimensional Hilbert space, for some* $d = 1, 2, \ldots$. *Let* $\mathcal{I}$ *be the set of isomorphism classes of AF* C^**-algebras of the form* $\mathfrak{M}_1/\mathfrak{O}(\mathfrak{L})$, *for* $\mathfrak{L}$ *an arbitrary maximal ideal of* $\mathfrak{M}_1$ *with* $\mathfrak{M}_1/\mathfrak{L} \cong \mathfrak{M}_1/\mathfrak{I}$. *Then* $\mathcal{I}$ *has exactly* $\max(1, \phi(d)/2)$ *elements.*

Proof. (i) A routine verification shows that $K_0(\mathfrak{I})$ is the set of all $f \in \mathcal{F}_1$ vanishing on θ. Since each linear piece of f has integer coefficients and θ is irrational, then f automatically vanishes on an open neighborhood of θ. So

$$\mathcal{F}_1/\mathfrak{m}_\theta = \mathcal{F}_1/\mathfrak{o}(\mathfrak{m}_\theta).$$

Recalling (10)-(12), we conclude

$$\mathfrak{M}_1/\mathfrak{I} \cong \mathfrak{M}_1/\mathfrak{O}(\mathfrak{I}).$$

(ii) Let G be the underlying group of $K_0(\mathfrak{M}_1/\mathfrak{I})$. Let d be the denominator of the smallest positive nonzero rational element of G. From the complete description of $\mu(\mathfrak{M}_1)$ given in [2] and [14] it follows that G has the form $\mathbb{Z}\frac{1}{d}$, for some integer $d > 0$. An elementary calculation of K_0 shows that $\mathfrak{M}_1/\mathfrak{I}$ is the C*-algebra of linear operators on d-dimensional Hilbert space. The rest now follows from Corollary 6.4. □

Problem. Prove or disprove: For all $n = 1, 2, 3, \ldots$ and $x, y \in \mathbb{R}^n$, $\mathrm{orb}(x) = \mathrm{orb}(y)$ iff $(G_x, c_x) = (G_y, c_y)$.

References

[1] A. Bigard, K. Keimel, and S. Wolfenstein, *Groupes et Anneaux Réticulés,* Lecture Notes in Mathematics, Springer-Verlag, Berlin, volume 608, 1971.

[2] F. Boca, An AF algebra associated with the Farey tessellation, *Canadian J. Mathematics*, 60 (2008) 975–1000.

[3] J.S.Dani, Density properties of orbits under discrete groups, *Journal of the Indian Mathematical Society*, 39 (1975) 189–218.

[4] C.Eckhardt, A noncommutative Gauss map, *Mathematica Scandinavica,* 108 (2011) 233–250.

[5] E. G. Effros, *Dimensions and C*-algebras*, CBMS Regional Conf. Series in Math., Vol. 46, Amer. Math. Soc., Providence, R.I., 1981.

[6] G. Ewald, *Combinatorial convexity and algebraic geometry*, Springer-Verlag, New York, 1996.

[7] K. R. Goodearl, *Notes on Real and Complex C*-Algebras*, Shiva Math. Series, Vol. 5, Birkhäuser, Boston, 1982.

[8] A. Guilloux, A brief remark on orbits of SL(2; $\mathbb{Z}$) in the Euclidean plane, *Ergodic Theory and Dynamical Systems,* 30 (2010) 1101–1109.

[9] G.H.Hardy, E.M.Wright, *An introduction to the theory of numbers,* Fifth Edition, Clarendon Press, Oxford, 1979.

[10] M. Laurent, A. Nogueira, Approximation to points in the plane by SL(2,$\mathbb{Z}$)-orbits, arXiv:1004.1326v1, 8 Apr 2010

[11] D. Mundici, Interpretation of AF C^*-algebras in Łukasiewicz sentential calculus, *Journal of Functional Analysis*, 65 (1986) 15–63.

[12] D. Mundici, Farey stellar subdivisions, ultrasimplicial groups, and K_0 of AF C^*-algebras, *Advances in Mathematics,* 68 (1988) 23-39.

[13] D.Mundici, The Haar theorem for lattice-ordered abelian groups with order-unit, *Discrete and Continuous Dynamical Systems*, 21 (2008) 537-549.

[14] D. Mundici, Revisiting the Farey AF algebra, *Milan Journal of Mathematics,* 79 (2011) 643–656.

[15] A. Nogueira, Orbit distribution on $\mathbb{R}^2$ under the natural action of SL$(2, \mathbb{Z})$, *Indag. Math.,* (N.S.), 13 (2002) 103–124.

[16] A. Nogueira, Lattice orbit distribution on $\mathbb{R}^2$, *Ergodic Theory Dynamical Systems,* 30 (2010) 1201–1214. Erratum, ibid., p. 1215.

[17] E.Witten, $SL(2,\mathbb{Z})$ action on three-dimensional conformal field theories with abelian symmetry, In: *From Fields to Strings: Circumnavigating Theoretical Physics*, Ian Kogan Memorial Collection (in 3 volumes), M. Shifman et al., Eds., World Scientific, Singapore, (2005), pp. 1173–1200.

(D. Mundici) Department of Mathematics "Ulisse Dini", University of Florence, Viale Morgagni 67/A, I-50134 Florence, Italy

E-mail address: mundici@math.unifi.it